

Role of Tannins-containing Shrub Legume *Calliandra calothyrsus* on Rumen Microbes Metabolisms


Firman Nasiu, Indonesia

Calliandra (*Calliandra calothyrsus*) is one of the most valuable forages for livestock such as cattle, sheep, goat, and buffalo.

INTRODUCTION



In Indonesia, calliandra is commonly used as the protein source feedstuff for livestock but it contains a chemical compound called **tannins** which, at excessive amount, can decrease nutritive value of feed produced in rumen.



Rumen is a vital compartment in stomach **which** contains millions of **microbes** including bacteria, fungi, and protozoa to support the animal in digesting feedstuff. But, these microorganisms can be disrupted by **tannins**.


OBJECTIVE

The aim of this study is to evaluate the effect of tannins compound on rumen microbial metabolisms which can be estimated from fermentation parameters including production of VFA (Volatile Fatty Acid), NH_3 , and protein microbial in rumen.

LITERATURE REVIEW

One of the good quality legume shrubs to provide adequate protein supply for ruminant is calliandra (*Calliandra calothyrsus*) (Ahn *et al.*, 1989). However, tannins in the plant can eliminate substrate availability to rumen microorganisms (McSweeney *et al.*, 2001), and decrease the rumen microbial metabolism to produce valuable nutrient for the animal (Tiemann *et al.*, 2008; Ørskov, 1992; Rimbawanto *et al.*, 2015; Bergmann, 1990).


Fermentation parameters value of calliandra analyzed by using *in vitro* technique.



Rimbawanto *et al.* (2015)

METHODOLOGY

In vitro technique following procedure by Tilley and Terry (1963) was applied to analyze rumen microbial metabolisms by estimating the value of VFA production, NH_3 , and protein microbes.



CONCLUSION

It is expected that there would be more data resulted from this laboratory experiment as the basic knowledge for further study of role of tannins on the animal performance including body weight gain and meat and milk production.

REFERENCES

- Ahn, J.H., Robertson, B.M., Elliott, R. Gutteridge, R.C. and Ford, C.W. (1989). Quality assessment of tropical browse legume: tannin content and protein degradation. *Animal Feed Science and Technology* 27, 147-156
- Bergman, E.N. (1990). Energy contribution of volatile fatty acids from the gastrointestinal tract in various species. *Physiol. Rev.* 70:567-590.
- *. D. (2017, September 25). *Calliandra calothyrsus* – Red Calliandra. Garden Manage. <https://es.gardenmanage.com/statuses/1000153292.html>.
- Lee, P. D. E. K. *Rumen fluid extraction made easy*. Progressive Dairy. <https://www.progressivedairy.com/topics/herd-health/rumen-fluid-extraction-made-easy>.
- McSweeney, C.S., B. Palmer, R. Bunch, and D.O. Krause. (2001). Effect of the tropical forage calliandra on microbial protein synthesis and ecology in the rumen. *Journal of Applied Microbiology* 90, 78-88.
- Ørskov, E.R. (1992). Protein Nutrition in Ruminants. 2nd Edition. Academic Press. New York, UEA.
- Plummer, D.T. (1987). An Introduction to Practical Biochemistry. McGraw Hill Ltd. Bombay, New Delhi. 352 pages.
- RAMM: Modeling Rumen Fermentation. Diamond V Knowledge Hub. <http://www.diamondv.com/blog/2017/08/10/ramm-modeling-rumen-fermentation/>.
- Ribeiro, L., Machado, F., Campos, M., Guimaraes, R., Tomich, T., Reis, L., & Coombs, C. (2015). Enteric methane mitigation strategies in ruminants: a review. *Revista Colombiana De Ciencias Pecuaria*, 28(2). <https://doi.org/10.17533/udea.rccpv28n2a02>
- Rimbawanto, E.A., L.M. Yusiati, E. Baliarti, R. Utomo. (2015). Effect of condensed tannin of leucaena and calliandra leaves in protein trash fish silage on *in vitro* ruminal fermentation, microbial protein synthesis and digestibility.
- Rose, A. L. (1970, January 1). *Digestive Physiology and Anatomy of Cows*. Vet in Training. <http://vetstudentresearch.blogspot.com/2015/05/digestive-physiology-and-anatomy-of-cows.html>.
- Tiemann, T.T., C.E. Lascano, H.-R. Wettstein, A.C. Mayer, M. Kreuzer, and H.D. Hess. (2008). Effect of the tropical tannin-rich shrub legumes *Calliandra calothyrsus* and *Flemingia macrophylla* on methane emission and nitrogen and energy balance in growing lambs. *Animal* 2:5, 790-799.
- Tilley, J.M.A. and R.A. Terry. (1963). A two-stage technique for the *in vitro* digestion of forage crop. *Journal of The British Grassland Society*. 18(2):104.
- Wang, N., Tan, F., Zhao, Y., Tsoi, C. C., Fan, X., Yu, W., & Zhang, X. (2016). Optofluidic UV-Vis spectrophotometer for online monitoring of photocatalytic reactions. *Scientific Reports*, 6(1). <https://doi.org/10.1038/srep28928>